



Neeuro Presents

# Improving Cognitive Health with Gamification and EEG

2022/2023 White Paper

# RESEARCH ON COGNITIVE TRAINING

The brain has been found to have the ability to control many aspects of functional thinking and this includes planning, organising, making rational decisions, remembering tasks and so much more. Our performance in these cognitive abilities can indicate how well we do our everyday tasks and whether we can live independently.

However, progressive changes occur in both physical and mental aspects as a person gets older [1-2] with cognitive decline occurring as early as in the thirties [94]. Certain parts of the brain shrink and can affect learning and other complex mental activities [95] the synapse structure (communication between neurons) may no longer be as effective [96], and the blood flow in the brain may decrease or be slower [97].

Nevertheless, slowing or preventing the onset of cognitive decline is possible by adopting a healthy lifestyle and cognitive interventions such as mental stimulation and cognitive training [98]. This is also supported by the National Institute of Health (NIH) which conducted a meta-study based on 250 high-quality publications and found cognitive training to be a highly effective protective factor in preventing cognitive decline [29].

Cognitive training has demonstrated benefits across all age groups, from young children to the elderly [33-44]. It has been

shown to play a positive role in improving fluid and crystallised intelligence [3-7] while also being sustainable [8-10] and transferable to daily functions [11-26]. At the same time, it has also been shown to be an effective non-pharmaceutical solution for people experiencing mental health issues [45-64].

With medical advancements, functional Magnetic Resonance Imaging (fMRI) and neuroimaging techniques have allowed researchers to observe the direct impacts cognitive training has on specific areas of the brain. These advancements have also given rise to the field of neuroplasticity (how malleable the brain is adapting to new stimulation), giving credence to the notion that our lives can be improved through training and stimulation of our brains [30-32]. Overall, the evidence found for the effectiveness of cognitive training has a strong scientific basis for combating cognitive decline.



# GAMES FOR THE BRAIN



Engaging in stimulating and purposeful activities is crucial for people to improve their quality of life. For those who have physical constraints or limited activity opportunities, digital games can offer a means for them to continue being engaged and mentally active. Driven by the National Science Foundation (NSF), neuroscientists and entertainment experts have initiated collaborations [68] to develop digital games to get a deeper understanding of brain functions and to provide new tools to boost attention and well-being. An example would be computer-based

cognitive training which is found to improve memory and performance of everyday tasks for seniors [66-67].

Moving forward, the health and technology industry is on its way to bringing more therapeutic games into the market [69], potentially creating a larger impact on those pursuing cognitive health and wellness.



# EEG-BASED COGNITIVE TRAINING


The use of electroencephalogram (EEG) by researchers in digital games is a recent advancement in the field of neuroscience. Numerous studies have investigated the impact of EEG in digital games and have found it to be effective in improving participants' cognitive abilities [1]. There are many associations between EEG biomarkers and cognitive functioning. For instance, there is a high correlation between the activation of theta brain wave activity (4–8 Hz) with one's working memory [70–71] while beta brain wave activity (14–30 Hz) is generally associated with one's attention level [72–73]. For tasks that require visuospatial abilities, participants who activate their alpha brain wave activity (9–13 Hz) performed better on mental rotations after going through EEG-based cognitive training [77]. Engaging individuals in tasks that induce these brain waves can help to dramatically improve their cognitive performance [75].

There may be some doubts concerning the possibility of diminishing cognitive gains during training due to cognitive overload, but this can be overcome with the use of an adaptive real-time monitoring system based on EEG biomarkers. By doing so, mental fatigue can be measured and effectively controlled hence allowing for participants to have a customised experience during gameplay to perform their best [74].

All in all, the use of EEG technology in facilitating cognitive training provides real-time brain wave measurements and gives accurate indications of a participant's mental state. The adaptive component found in the cognitive training programme also enhances the experience hence effectively reaping maximum benefits.

# NEEURO MEMORIE GAMES

Neeuro's Memorie mobile application equipped with Neeuro's EEG headband, SenzeBand 2 offers a unique solution for brain training, which can be done in a fun manner, anytime and anywhere.







# SUSHI RECALL


Sushi Recall is a fun version of n-back task that is most commonly used in cognitive assessments to measure working memory [84]. Using the n-back task has also been shown to increase both brain thickness and surface areas which indicates better performance in complex cognitive tasks such as language comprehension, learning, and reasoning [85].

During gameplay, the player has to remember a variety of sushi types supplied by

a conveyor belt. As one or more sushi disappears, the player has to recall the last sushi. The number of sushi disappearing will progressively increase to make the task more difficult as the player moves to the next level. Hence, for players to succeed in each level, they will need to continuously activate and manipulate information from their working memory.

The objective of sushi recall is to essentially allow players to practice temporarily retaining

pieces of information that can be vital to conduct a variety of everyday tasks. This ability to hold information allows for better everyday reasoning and decision making such as dialling a new telephone number or recalling where you might have placed your items such as a book or a mobile phone.



# DOT CONNECT

Dot Connect is inspired by mental rotation tasks [86] where the player will have to activate their spatial abilities by identifying a given 2D line segment after which they will have to replicate, flip and rotate a separate line segment as per instructions from the game. While seldom discussed compared to short-term memory loss when it comes to cognitive decline, our spatial abilities are an important subset of our general intelligence [87]. These cognitive skills are pervasive

in various everyday tasks, such as reading maps, driving [88], and during mathematical performance [89].

In the beginner levels of Dot Connect, the player will only need to mentally rotate the board in 90 and 180 degrees clockwise or anticlockwise. A colour guide is provided in the earlier stages to familiarise players with the rules of the game, after which it will be removed as players progress into later stages and get more competent.

Additionally, the sensors from our Senzeband will allow evaluation of players' visual spatial attention and focus [90] hence providing accurate measurements in tracking their overall progress.



# MINDCOPTER

Mind Copter leverages studies that show the association of EEG biomarkers to facilitate attention training [72, 73]. The Senzeband captures real-time measurement of the player's attention level which not only provides an extra dimension to game scoring but also tailors an adaptive training programme to improve their selective and divided attention capabilities. These cognitive skills are important to manage distraction so that unimportant details can be

tuned out and one can finally focus on what matters [91]. This will also be useful to help concentration during a task or event for a prolonged period.

The player will take on the role of a pilot and will need to activate their attention to airborne the helicopter and perform rescue missions for stranded survivors. The player will need to sustain their attention to drop off care packages for survivors and to wilfully release attention so that the helicopter can

descend and rescue them. As the game progresses, the helicopter gets 'heavier' and harder to control with more obstacles introduced.





# PYRAMID SOLITAIRE

Pyramid Solitaire is a modified version of the classic Solitaire game in which gameplay has been demonstrated to improve cognitive functions such as short-term memory [78]. Leveraging on the concept of neuroplasticity, when players have to frequently activate their working memory to recall the value of the cards [79], the brain will create or strengthen new neural pathways and hence have the potential to remain robust even in the cases of Alzheimer's Disease

(AD) [82].

The game requires players to match two cards that will add up to the value of 13. Players will also activate their decision-making skills as they decide which strategy or pathway to take first, depending on the cards they have on hand and the cards available in the pyramid. With the right strategical decision, they will not chance upon a situation where they unlock a tier that has no other cards to match. Unlike classic pyramid

Solitaire where unused cards are promptly disposed of, Pyramid Solitaire allows players to stash their cards for later use. However, cards are faced downwards so they will need to remember the values of the cards that are stowed away. This requires memory work and any wrong guesses will lead to a score penalty.



# MULTITASK MASTER

Multitasker Master was created on ideas of task-switching paradigms [92] and the drive towards building cognitive flexibility that helps people pursue complex tasks such as efficient problem-solving and creativity [93].

The game is uniquely split into two screens with tasks to complete on both sides. Points are granted when tasks are completed screen by screen. This means the player has to complete screen 1, then move on to screen 2

then back to screen 1. The tasks to complete are different in nature hence encouraging the player to use various cognitive functions quickly and efficiently. For example, screen 1 could be displaying a mathematical sum that requires mental calculation while screen 2 displays a pattern recognition problem that requires the player to identify and analyse patterns between figures. As the player progresses, the time given for the player to solve questions on each screen reduces.



# CONCLUSION

We are in a world of rapid information growth and experiencing the age of attention crisis. With unprecedentedly longer life spans, people are experiencing a cognitive decline earlier than expected. These are the factors that place a great challenge for us to maintain cognitive fitness throughout life. To enhance our cognitive well-being and prevent cognitive decline, it's important to engage in ample of mental exercises in addition to a healthy lifestyle.

The effectiveness of brain training has been advocated by decades of scientific research. While mobile applications and computer games become increasingly inevitable in today's world, so is the convergence of health, technology and games. Mobile games paired with lightweight EEG sensors make it possible to facilitate closed-loop cognitive training in daily life, offering adaptive, handy and powerful training programmes for effective brain exercise for everyone, anytime and anywhere.



# ABOUT NEEURO

+65 6397 5153

contact@neeuro.com

www.neeuro.com

Neeuro is a global company that specialises in utilising Brain-Computer Interface technology to maximise the potential of users' neurological agility and fitness. Since its inception in 2013, Neeuro's extensive portfolio and an array of digital therapeutic and brain fitness solutions are backed by clinically validated research by A\*STAR, an institution widely known as being at the forefront of Singapore's research endeavours.

Its core technology, NeeuroOS, is a platform that empowers healthcare professionals, researchers, and third-party developers with an Artificial Intelligence (AI) driven ability to analyse the brain signals of users, measuring mental states including but not limited to attention, relaxation, mental workload, and fatigue. Neeuro's holistic offerings numerous potential avenues to explore complementary mental health options for children with ADHD, patients with stroke, cognitive rehabilitation, and many other neurological challenges.

# REFERENCES

## **LINEAR TREND OF COGNITIVE DECLINE WHILE AGING**

1. Tucker-Drob EM & Salthouse TA. Adult age trends in the relations among cognitive abilities. *Psychol. Aging* 23, 453–460 (2008).
2. Anguera JA, Boccanfuso J, Rintoul JL et al. Video game training enhances cognitive control in older adults. *Nature*. 2013 Sep 5; 501(7465):97–101.

## **COGNITIVE TRAINING FOR FLUID INTELLIGENCE**

3. Jaeggi SM, Buschkuhl M, Jonides J & Perrig WJ. Improving fluid intelligence with training on working memory. *Proc Natl Acad Sci USA*. 105, 6829–6833 (2008).
4. Zhao X, Wang YX, et al., Effect of updating training on fluid intelligence in children. *Chin Sci Bull*. 2011 Jul; 56 (21): 2202–5.
5. Bergman NS, Soderqvist S, et al., Gains in fluid intelligence after training non-verbal reasoning in 4-year-old children: A controlled, randomized study. *Dev Sci*. 2011 May; 14 (3): 591–601.
6. Rudebeck SR, Bor D, et al., A potential spatial working memory training task to improve both episodic memory and fluid intelligence. *PLoS One*. 2012; 7 (11): e50431.

## **COGNITIVE TRAINING FOR CRYSTAL INTELLIGENCE**

7. Alloway TP, Alloway RG, The efficacy of working memory training in improving crystallized intelligence. *Nature Proceedings*. 2009 Sep.

## **COGNITIVE TRAINING IS SUSTAINABLE**

8. Bennett SJ, Holmes J, et al., Computerized memory training leads to sustained improvement in visuospatial short-term memory skills in children with Down syndrome. *Am J Intel Dev Disab*. 2013; 118 (3): 179–92.
9. Brehmer Y, Westerberg H, et al., Working-memory training in younger and older adults: Training gains, transfer, and maintenance. *Front Hum Neurosci*. 2012 Mar; 6: 63.
10. Wolf D, Fischer FU, et al., Structural integrity of the corpus callosum predicts long-term transfer of fluid intelligence-related training gains in normal aging. *Hum Brain Mapp*. 2012 Sep 11.

## **COGNITIVE TRAINING IS TRANSFERABLE FOR DAILY TASKS**

11. Chein JM, Morrison AB, Expanding the mind's workspace: Training and transfer effects with a complex working memory span task. *Psychon Bull Rev*. 2010 Apr; 17 (2): 193–99.
12. Heinzl S, Schulte S, et al., Working memory training improvements and gains in non-trained cognitive tasks in young and older adults. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn*. 2013 May 2
13. Holmes J, Gathercole SE, et al., Adaptive training leads to sustained enhancement of poor working memory in children. *Dev Sci*. 2009 Jul; 12 (4): F9–15.
14. Kundu B, Sutterer DW, et al., Strengthened effective connectivity underlies transfer of working memory training to tests of short-term memory and attention. *J Neurosci*. 15 May 2013; 33 (20): 8705–15.
15. Salminen T, Strobach T, et al., On the impacts of working memory training on executive functioning. *Front Hum Neurosci*. 2012; 6: 166.
16. Loosli SV, Buschkuhl M, et al., Working memory training improves reading processes in typically developing children. *Child Neuropsychol*. 2012; 18 (1): 62–78.
17. Schweizer S, Hampshire A, et al., Extending brain-training to the affective domain: Increasing cognitive and affective executive control through emotional working memory training. *PLoS One*. 2011; 6 (9): e24372.
18. Von Bastian CC, Oberauer K, Distinct transfer effects of training different facets of working memory capacity. *J Mem Lang*. 2013 Jul; 69 (1): 36–58.
19. Lilienthal L, Tamez E, et al., Dual n-back training increases the capacity of the focus of attention. *Psychon Bull Rev*. 2013 Feb; 20 (1): 135–41.
20. Minear M, Shah P, Training and transfer effects in task switching. *Mem Cognit*. 2008 Dec; 36 (8): 1470–83.
21. Schmiedek F, Lövdén M, et al., Hundred days of cognitive training enhance broad cognitive abilities in adulthood: Findings from the COGITO study. *Front Aging Neurosci*. 2010 Jul 13; 2.
22. Thorell LB, Lindqvist S, et al., Training and transfer effects of executive functions in preschool children. *Dev Sci*. 2009 Jan; 12 (1): 106–13.
23. Rueda MR, Checa P, et al., Enhanced efficiency of the executive attention network after training in preschool children: Immediate and effects after two months. *Dev Cogn Neurosci*. 2012 Feb 15; 2 Suppl 1: S192–204.
24. Jaeggi SM, Studer-Luethi B, et al., The relationship between n-back performance and matrix reasoning—implications for training and transfer. *Intelligence*. 2010; 38: 625–35.
25. Jaeggi SM, Buschkuhl M, et al., Short- and long-term benefits of cognitive training. *PNAS*. 2011 Jun 21; 108 (25): 10081–86.
26. Jausovec N, Jausovec K, Working memory training: Improving intelligence—changing brain activity. *Brain Cogn*. 2012; 79: 96–106.

## COGNITIVE ASSESSMENT

27. Randolph C, Tierney MC, Mohr E, Chase TN. The Repeatable Battery for the Assessment of Neuro-psychological Status (RBANS): preliminary clinical validity. *J Clin Exp Neuropsychol*. 1998 20(3): 310–9.
28. Barbara Bradley Hagerty, Forget About It: Your Middle-Aged Brain Is Not On the Decline. NPR, Mar 15, 2016.

## LARGE-SCALE META STUDY

29. John W Williams, Brenda L Plassman, James Burke, Tracey Holsinger and Sophiya Benjamin. Preventing Alzheimer's Disease and Cognitive Decline. Evidence Reports: AHRQ publication (2010).

## MRI VERIFIED BRAIN CHANGES - NEUROPLASTICITY

30. Olesen PJ, Westerberg H, Klingberg T. Increased prefrontal and parietal activity after training of working memory. *Nat Neurosci*. 2004 Jan; 7 (1): 75–79.
31. Schweizer S, Grahm J, et al., Training the emotional brain: Improving affective control through emotional working memory training. *J Neurosci*. 2013 Mar 20; 33 (12): 5301–11.
32. Vartaniana O, Jobidona ME, Working memory training is associated with lower prefrontal cortex activation in a divergent thinking task. *Neuroscience*. 2013 Apr 16; 236: 186–94.

## COGNITIVE TRAINING FOR CHILDREN

33. Mackey AP, Hill SS, et al., Differential effects of reasoning and speed training in children. *Dev Sci*. 2011 May; 14 (3): 582–90.
- Cognitive Training for the Elderly
34. Buschkuhl M, Jaeggi SM, et al., Impact of working memory training on memory performance in old-old adults. *Psychol Aging*. 2008 Dec; 23 (4): 745–53.
35. Li SC, Schmiedek F, et al., Working memory plasticity in old age: Practice gain, transfer, and maintenance. *Psychol Aging*. 2008 Dec; 23 (4): 731–42.
36. Wolinsky FD, Vander Weg MW, et al., A randomized controlled trial of cognitive training using a visual speed of processing intervention in middle aged and older adults. *PLoS One*. 2013 May 1; 8 (5): e61624.
37. Zinke K, Zeintl M, Working memory training and transfer in older adults: Effects of age, baseline performance, and training gains. *Dev Psychol*. 2013 May 20 .
38. Liu-Ambrose T, Nagamatsu LS, et al., Resistance training and executive functions: A 12-month randomized controlled trial. *Arch Intern Med*. 2010; 170 (2): 170–78.
39. von Bastian CC, Langer N, et al., Effects of working memory training in young and old adults. *Mem Cognit*. 2013 May; 41 (4): 611–24.
40. Richmond LL, Morrison AB, et al., Working memory training and transfer in older adults. *Psychol Aging*. 2011 Dec; 26 (4): 813–22.
41. Basak C, Boot WR, et al., Can training in real-time strategy video game attenuate cognitive decline in older adults? *Psychol Aging*. 2008; 23 (4): 765–77.
42. Carretti B, Borella E, et al., Gains in language comprehension relating to working memory training in healthy older adults. *Int J Geriatr Psychiatry*. 2013 May; 28 (5): 539–46.
43. Borella E, Carretti B, et al., Working memory training in older adults: Evidence of transfer and maintenance effects. *Psychol Aging*. 2010; 25 (4): 767–778.
44. Ball K, Berch DB, et al. Effects of cognitive training interventions with older adults: A randomized controlled trial. *JAMA*. 2002 Nov 13; 288 (18): 2271–81.

## COGNITIVE THERAPY

45. Boron JB, Willis SL, et al., Cognitive training gains as a predictor of mental status. *J Gerontol*. 2007 Jan; 62B (1): P45–51.
46. Bell M, Bryson G, et al., Cognitive remediation of working memory deficits: Durability of training effects in severely impaired and less severely impaired schizophrenia. *Acta Psychiatr Scand*. 2003; 108: 101–9.
47. Dahlin KIE, Effects of working memory training on reading in children with special needs. *Reading and Writing*. 2011; 24: 479–91.
48. Gray SA, Chaban P, et al., Effects of a computerized working memory training program on working memory, attention, and academics in adolescents with severe LD and comorbid ADHD: A randomized controlled trial. *J Child Psychol Psychiatry*. 2012 Dec; 53 (12): 1277–84.
49. Green CT, Long DL, et al., Will working memory training generalize to improve off-task behavior in children with attention-deficit/hyperactivity disorder? *Neurotherapeutics*. 2012 Jul; 9 (3): 639–48.
50. Hardy KK, Willard VW, et al., Working memory training in survivors of pediatric cancer: A randomized pilot study. *Psycho-Oncology*. 2012 Dec 2



51. Holmes J, Gathercole SE, et al., Working memory deficits can be overcome: Impacts of training and medication on working memory in children with ADHD. *Appl Cognit Psychol*. 2010 Sep; 24 (6): 827–36.
52. Houben K, Wiers RW, et al., Getting a grip on drinking behavior: Training working memory to reduce alcohol abuse. *Psychol Sci*. 2011 Jul; 22 (7): 968–75.
53. Kesler S, Hadi Hosseini SM, et al., Cognitive training for improving executive function in chemotherapy-treated breast cancer survivors. *Clin Breast Cancer*. 2013 Aug; 13 (4): 299–306.
54. Klingberg T, Fernell E, et al., Computerized training of working memory in children with ADHD—a randomized, controlled trial. *J Am Acad Child Adolesc Psychiatry*. 2005; 44 (2): 177–86.
55. Kray J, Karbach J, Can task-switching training enhance executive control functioning in children with attention deficit/-hyperactivity disorder? *Front Hum Neurosci*. 2011; 5: 180.
56. McGurk SR, Mueser KT, et al., Cognitive training and supported employment for persons with severe mental illness: One-year results from a randomized controlled trial. *Schiz Bull*. 2005; 31 (4): 898–909.
57. Nagamatsu LS, Handy TC, et al., Resistance training promotes cognitive and functional brain plasticity in seniors with probable mild cognitive impairment. *Arch Intern Med*. 2012 Apr 23; 172 (8): 666–68.
58. Owens M, Koster EH, et al., Improving attention control in dysphoria through cognitive training: Transfer effects on working memory capacity and filtering efficiency. *Psychophysiology*. 2013 Mar; 50 (3): 297–307.
59. Prins PJ, DAVIS S, et al., Does computerized working memory training with game elements enhance motivation and training efficacy in children with ADHD? *Cyberpsychol Behav Soc Netw*. 2011 Mar; 14 (3): 115–22.
60. Soderqvist S, Nutley SB, et al., Computerized training of non-verbal reasoning and working memory in children with intellectual disability. *Front Hum Neurosci*. 2012; 6: 271.
61. Subramaniam K, Luks TL, et al., Computerized cognitive training restores neural activity within the reality monitoring network in schizophrenia. *Neuron*. 2012 Feb 23; 73: 842–53.
62. Van der Molen MJ, Van Luit JE, et al., Effectiveness of a computerised working memory training in adolescents with mild to borderline intellectual disabilities. *J Intellect Disabil Res*. 2010 May; 54 (5): 433–47.
63. Roughan L, Hadwin JA, The impact of working memory training in young people with social, emotional and behavioral difficulties. *Learning and Individual Differences*. 2011 Dec; 21 (6): 759–64.
64. Neville HJ, Stevens C, et al., Family-based training program improves brain function, cognition, and behavior in lower socioeconomic status preschoolers. *PNAS*. 2013 Jul 1, 110(29), p12138–12143

#### **TRAINING IN DIFFERENT AGE GROUPS**

65. Karbach J, Kray J, How useful is executive control training? Age differences in near and far transfer of task-switching training. *Dev Sci*. 2009 Nov; 12 (6): 978–90.

#### **TRAINING GAMES**

66. Sunghee H. Tak, Cornelia Beck, and Song Hee Hong. Feasibility of providing computer activities for nursing home residents with dementia. *Nonpharmacol Ther Dement*. 2013; 3(1): 1–10.
67. Anne Corbett et al. The Effect of an Online Cognitive Training Package in Healthy Older Adults: An Online Randomized Controlled Trial. *JAMDA* 16 (2015) 990e997
68. Workshop on Interactive Media, Attention and Well-Being. <http://go.nature.com/t9mvqc>
69. Daphne Bavelier & Richard J. Davidson. Brain training: Games to do you good. *Nature* 494, 425–426 (28 February 2013)

#### **EEG BASED TRAINING**

70. Kahana, M.J., Sekuler, R., Caplan, J.B., Kirschen, M., Madsen, J.R., 1999. Human theta oscillations exhibit task dependence during virtual maze navigation. *Nature* 399, 781–784.
71. Klimesch, W., Doppelmayr, M., Stadler, W., Pollhuber, D., Sauseng, P., Rohm, D., 2001. Episodic retrieval is reflected by a process specific increase in human electroencephalographic theta activity. *Neurosci. Lett.* 302, 49–52
72. Egner, T., Gruzelier, J.H., 2001. Learned self-regulation of EEG frequency components affects attention and event-related brain potentials in humans. *Neuroreport* 12, 4155–4159.
73. Rossiter, T.R., LaVaque, T.J., 1995. A comparison of EEG biofeedback and psychostimulants in treating attention deficit hyperactivity disorders. *J. Neurother.* 48–59
74. Baldwin, C. L., and Penaranda, B. N. (2012). Adaptive training using an artificial neural network and EEG metrics for within- and cross-task workload classification. *Neuroimage* 59, 48–56.
75. Thomas KP, Vinod AP, Guan C. Enhancement of attention and cognitive skills using EEG based neurofeedback game. *Neural Engineering (NER)*, 2013 6th International IEEE/EMBS Conference on. P 21 – 24
76. Vernon D, Egner T, Cooper N, Compton T, Neilands C, Sheri A, Gruzelier J. The effect of training distinct neurofeedback protocols on aspects of cognitive performance. *Int J Psychophysiol*. 2003 Jan;47(1):75–85.

77. Hanslmayr, S., Sauseng, P., Doppelmayr, M., Schabus, M., Klimesch, W., 2005. Increasing individual upper alpha power by neurofeedback improves cognitive performance in human subjects. *Appl. Psychophysiol. Biofeedback* 30, 1–10.

## COGNITIVE (OTHERS)

78. Shinya Uchida & Ryuta Kawashima. Reading and solving arithmetic problems improves cognitive functions of normal aged people: a randomized controlled study. *AGE* (2008) 30:21–29

79. Gerd Kempermann, Daniela Gast and Fred H. Gage. Neuroplasticity in old age: Sustained fivefold induction of hippocampal neurogenesis by long-term environmental enrichment. *Annals of Neurology*, Vol 52, Issue 2, pages 135–143, August 2002

80. Buitenweg J, Murre JM, Ridderinkhof KR. Brain training in progress: a review of trainability in healthy seniors. *Front Hum Neurosci.* 2012 Jun 21;6:183.

81. Nelson Cowan, The Magical Mystery Four: How is Working Memory Capacity Limited, and Why? *Curr Dir Psychol Sci.* 2010 Feb 1; 19(1): 51–57.

82. Sergio Machado, Marlo Cunha et al. ALZHEIMER'S DISEASE AND IMPLICIT MEMORY. *Arq Neuropsiquiatr* 2009;67(2-A):334-342 334

83. Rui Nouchi et al. Brain Training Game Improves Executive Functions and Processing Speed in the Elderly: A Randomized Controlled Trial. *PlosOne*, Volume 7, Issue 1, e29676.

84. RuGazzaniga, Michael S.; Ivry, Richard B.; Mangun, George R. (2009). *Cognitive Neuroscience: The Biology of the Mind* (2nd ed.).

85. Francisco J. Román, Lindsay B. Lewis et. al. Gray matter responsiveness to adaptive working memory training: a surface-based morphometry study. *Brain Structure and Function*, pp 1–14, 2015

86. Vandenberg, S. G., & Kuse, A. R. (1978, December). Mental rotations, a group test of three-dimensional spatial ...Retrieved November 1, 2022, from <https://journals.sagepub.com/doi/abs/10.2466/pms.1978.47.2.599>

87. Howard Gardner, *Frames of Mind: The Theory of Multiple Intelligences*, 1983.

88. Moen, K.C. et al. (2020) Strengthening spatial reasoning: Elucidating the attentional and neural mechanisms associated with mental rotation skill development - cognitive research: Principles and implications, SpringerOpen. Springer International Publishing. Available at: <https://cognitiveresearchjournal.springeropen.com/articles/10.1186/s41235-020-00211-y> (Accessed: November 2, 2022).

89. Mix K. S., Levine S. C., Cheng Y. L., Young C., Hambrick D. Z., Ping R., Konstantopoulos S. (2016). Separate but correlated: The latent structure of space and mathematics across development. *Journal of Experimental Psychology: General*, 145(9), 1206–1227. <https://doi.org/10.1037/xge0000182>

90. *The Oxford Handbook of Attention*, Edited by Anna C. Nobre and Sabine Kastner, 2014.

91. Stevens, C. and Bavelier, D. (2012) The role of selective attention on academic foundations: A cognitive neuroscience perspective, *Developmental cognitive neuroscience*. Elsevier. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3375497/> (Accessed: November 2, 2022)

92. Rogers, R. and Monsell, S. (no date) Costs of a predictable switch between simple cognitive tasks - researchgate, Research Gate. Available at: [https://www.researchgate.net/publication/232496441\\_Costs\\_of\\_a\\_Predictable\\_Switch\\_Between\\_Simple\\_Cognitive\\_Tasks](https://www.researchgate.net/publication/232496441_Costs_of_a_Predictable_Switch_Between_Simple_Cognitive_Tasks) (Accessed: November 2, 2022).

93. Ionescu, T. (2011) Exploring the nature of cognitive flexibility, *New Ideas in Psychology*. Pergamon. Available at: <https://www.sciencedirect.com/science/article/abs/pii/S0732118X11000705> (Accessed: November 2, 2022).

94. Gonzales, M. et al. (2022) "Biological aging processes underlying cognitive decline and neurodegenerative disease," *Journal of Clinical Investigation*, 132(10). Available at: <https://doi.org/10.1172/jci158453>.

95. Peters, R. (2006) Ageing and the brain, *Postgraduate medical journal*. BMJ Group. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2596698/> (Accessed: November 7, 2022).

96. Petralia - Petralia, R.S., Mattson, M.P. and Yao, P.J. (2014) Communication breakdown: The impact of ageing on Synapse Structure, *Ageing research reviews*. U.S. National Library of Medicine. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4094371/> (Accessed: November 7, 2022).

97. Juttukonda - Characterizing cerebral hemodynamics across the adult lifespan with arterial spin labeling MRI data from the Human Connectome Project-Aging

Juttukonda, M. R., Li, B., Almaktoom, R., Stephens, K. A., Yochim, K. M., Yacoub, E., Buckner, R. L., & Salat, D. H. (2021). *NeuroImage*, 230, 117807. <https://doi.org/10.1016/j.neuroimage.2021.117807>

98. Risk reduction of cognitive decline and dementia: Who guidelines (2019) World Health Organization. World Health Organization. Available at: <https://apps.who.int/iris/handle/10665/312180> (Accessed: November 7, 2022).